

## Points to Consider

- ✓ Children's Medical Group does not recommend the use of prescription strength treatments.
- ✓ Prescription lice treatment is not available from our providers.
- ✓ Over-the-counter treatments are the safest and most effective as long as directions are followed carefully.
- ✓ If you suspect head lice in a baby less than 2 months of age, please call Children's Medical Group at the number listed on our website.

## **Treating Headlice**

- Crème rinse, such as Nix (not shampoo) is typically more effective than shampoos. In addition to what is listed below, please follow the directions completely.
  - Before using Nix, clean hair with any shampoo; then, towel dry hair.
  - Continue massaging Nix crème rinse into hair for a <u>full 10 minutes</u>—any air pockets left (even the size of a sesame seed) may allow lice to survive.
  - Nix leaves a residue on the hair for up to 2 weeks.
- ✓ If you choose lice shampoo, such as RID or Lice Away, **please follow the directions carefully**.
  - Apply to **dry** hair, saturate ALL hair down to roots, add water, and lather.
  - Continue massaging shampoo into the hair for a <u>full 10 minutes</u>—any air pockets left (even the size of a sesame seed) may allow lice to survive.
- ✓ To destroy lice and eggs on bedding, furniture, carpet, clothing, etc.:
  - Do not use chemical sprays—they are not effective.
  - Vacuum thoroughly and discard the bag <u>or</u> wash the canister in hot, soapy water.
  - Wash all clothing in hot water.

## Follow-up Care

- Nits appear as tiny clear, whitish flakes that are tightly adhered to the shaft of individual hairs located at or up to 1-2 inches from the scalp.
- **OPTIONAL**: "Nit Picking" **MAY** help remove nits and eggs. If you choose this step:
  - It can be done every 2-3 days for 2-3 weeks following the initial crème or shampoo treatment.
  - Work with tiny sections of hair & follow a structured pattern, so as to cover all areas.
  - Search for no longer than 15-20 minutes at a time---rest for a while before restarting, otherwise eyes may 'tune-out' to the view of the tiny nits.
  - If you suspect a new outbreak of lice 7-10 days after treatment, please contact Children's Medical Group to see if an additional treatment is appropriate for your child.